Reduction of $SU_f(3) \supset SO(3) \supset A_4$-The scalar potential.

J. D. Vergados*
ARC Centre of Excellence in Particle Physics at the Terascale and Centre for the Subatomic Structure of Matter (CSSM), University of Adelaide, Adelaide SA 5005, Australia

We examine the possibility of the flavor symmetry being $SU_f(3)$ in which the phenomenologically successful discreet symmetry A_4 is embedded. The allowed quadratic and quartic potentials are constructed exploiting the full symmetry chain $SU(3) \supset SO(4) \supset A_4$. The special case of the $SU(3)$ decaplet and octet scalar fields is analyzed.

PACS numbers:

arXiv:1604.00678v1 [physics.gen-ph] 19 Mar 2016

INTRODUCTION

For three generations of fermions the maximum horizontal symmetry is expected to be $SU(3)$ which contains as a subgroup the phenomenologically acceptable discreet symmetry A_4. It is known the the irreducible representations of A_4 are three singlets indicated as $\underline{1}$, $\underline{1}^{'}$, $\underline{1}^{''}$ and a triplet indicated as $\underline{3}$. The irreducible representations of $SU(3)$ are specified by two non negative integers (λ, μ). These are related to the young tableaux $[f_1, f_2, f_3]$, $f_1 \geq f_2 \geq f_3$ of $U(3)$ by $\lambda = f_1 - f_2$ and $\mu = f_2 - f_3$ In particle physics the $SU(3)$ representations are indicated merely by their dimension. This is not adequate for our purposes since there may be more than one irreducible representations with the same dimension.(μ, λ) is the adjoined (conjugate) of (λ, μ). Representations of the type (λ, λ) are self-adjoined.

The $SO(3)$ or $R(3)$ group is well known angular momentum theory. The A_4 is a discreet group isomorphic to the tetrahedral symmetry that has recently become popular in particle physics in connection with the neutrino mass. It is obviously a subgroup of $SO(3)$. Since, however, there exist three quark and lepton families, one may suppose an $SU(3)$ flavor symmetry and would like to see the A_4 emerging as a subgroup of $SU(3)$.

THE STRUCTURE OF A_4

S_4 is the group permutations of 4 elements. It contains

1. The identity:

$$C_1 \Leftrightarrow = \begin{pmatrix} 1\,2\,3\,4 \\ 1\,2\,3\,4 \end{pmatrix} = (1\,1),\ (2\,2),\ (3\,3),\ (4\,4)$$

2. 6 transpositions

$$C_2 \Leftrightarrow (12),\ (1\,3),\ (1\,4),\ (2\,3),\ (2\,4)$$

3. 8 cyclic permutations of length 3

$$C_3 \Leftrightarrow (1\,2\,3),\ (3\,2\,1),\ (1\,2\,4),\ (4\,2\,1),\ (1\,3\,4),\ (4\,3\,1),\ (2\,3\,4),\ (4\,3\,2)$$

4. 6 cyclic permutations of length 4

$$C_4 \Leftrightarrow (1\,2\,3\,4),\ (1\,2\,4\,3),\ (1\,3\,2\,4),\ (1\,3\,4\,2),\ (1\,4\,2\,3),\ (1\,4\,3\,2)$$

5. 3 products of two transpositions

$$C_5 \Leftrightarrow (1\,2)(3\,4),\ (1\,3)(2\,4),\ (1\,4)(2\,3)$$

* Permanent address: University of Ioannina, Ioannina, Greece
email:vergados@uoi.gr

We note that the 12 elements of the classes C_1, C_3, C_5 form a group, ${}_4 = C_1 + C_3 + C_5$. The group ${}_4$ is isomorphic to the tetrahedral group.
with respect to A_4 the class C_3 splits into two classes:

$$C_3 \Leftrightarrow (1\,2\,3),\ (4\,2\,1),\ (4\,3\,2),\ (1\,3\,4), \quad C_3 \Leftrightarrow (1\,2\,4),\ (4\,2\,1),\ (4\,3\,1),\ (2\,3\,4)$$

So there exist 4 classes and, hence, 4 irreducible representations with dimensions constrained by:

$$\sum_{i=1}^{3} \ell_i^2 = 12 \tag{1}$$

which has a unique solution:

$$\ell_1 = 1,\ \ell_2 = 1,\ \ell_3 = 1,\ \ell_4 = 3 \tag{2}$$

The three dimensional is the most important. We have :

- The identity element, 3×3.
- The elements of class C_5 commute and they can be simultaneously diagonalized. Since they are orthogonal with determinant equal to one they can be cast into the form:

$$t_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad t_2 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad t_3 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

  This representation is often called real. The matrix t_2 is sometimes denoted by$S^{'}$.

- the operator $(1\,2\,3)$ induces the transformation $(x_1, x_2, x_3) \rightarrow (x_2, x_3, x_1)$ and $(1\,3\,2)$ the transformation $(x_1, x_2, x_3) \rightarrow (x_3, x_1, x_2)$. They are thus represented as:

$$(132) \Leftrightarrow a(0) = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad (123) \Leftrightarrow b(0) = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}.$$

  In terms of them we get

$$a(i) = t(i)a(0)t(i), \quad b(i) = t(i)b(0)t(i), \quad i = 1, 2, 3.$$

  In other words:

$$(1) = \begin{pmatrix} 0 & 0 & -1 \\ -1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, (2) = \begin{pmatrix} 0 & 0 & 1 \\ -1 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}, (3) = \begin{pmatrix} 0 & 0 & -1 \\ 1 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}$$

$$(1) = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{pmatrix}, (2) = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & -1 \\ 1 & 0 & 0 \end{pmatrix}, (3) = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & -1 \\ -1 & 0 & 0 \end{pmatrix}.$$

  The matrix $b(2)$ is sometimes denoted by $T^{'}$. The elements $T = (123)$ and $S = (14)(23)$ are the generators of the group.

  Thus one can select a basis in which T is diagonal. Indeed

$$T = U^{+}b(0)U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, A = \sqrt{3}U = \begin{pmatrix} 1 & 1 & 1 \\ \omega & 1 & \omega^2 \\ \omega^2 & 1 & \omega \end{pmatrix}, \omega = e^{\frac{2\pi i}{3}}$$

Then

$$U^+ t_2 U \to S = \frac{1}{3}\begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}$$

$$U^+ t_3 U \to S_3 = -1 + \frac{2}{3}\begin{pmatrix} 1 & \omega & \omega^2 \\ \omega^2 & 1 & \omega \\ \omega & \omega^2 & 1 \end{pmatrix}$$

$$U^+ t_1 U \to S_1 = -1 + \frac{2}{3}\begin{pmatrix} 1 & \omega^2 & \omega \\ \omega & 1 & \omega^2 \\ \omega^2 & \omega & 1 \end{pmatrix}$$

APPLICATION IN $SU(3) \supset SO(3) \supset A_4$ CHAIN

We start with $SU(3) \supset SO(3)$ basis. The states in this chain are specified by the $SU(3)$ quantum numbers (λ, μ) as well as the angular momentum quantum numbers (L, M). The construction of these states is well known [1]. In general an additional quantum number κ is needed,

$$|(\lambda, \mu)\kappa, L, M.$$

It is not easy to construct an operator, which is made up from the generators of $SU(3)$ and commutes with L^2 and L_z, with eigenvalues that specify κ. So, whenever this additional quantum number is needed to remove $L-$ degeneracy, we will use an algorithm already developed [1].

The reduction of the $SO(3)$ representations to A_4 has been considered in the past, with elegant but abstract and rather obscure for most physicists mathematical techniques [2–4]. We will try to do this in a more direct way motivated by physics. This way we will get a handle on how to construct the $SU(3) \supset SO(3) \supset A_4$ invariants and break the $SU(3)$ down to A_4

In order to realize the cyclic permutation (1 2 3) of A_4 in the angular momentum basis we need consider representations, which involve at least three particles. Then a basis is specified by the magnetic quantum numbers$\{m_1, m_2, m_3\}$ for the three spin one objects. Since the operator (1 2 3) commutes with $L_z = L_z(1) + L_z(2) + L_z(3)$, both of them can be simultaneously diagonalized. This means that it is adequate to consider each $m = m_1 + m_2 + m_3$ separately.

Each particle is put in the fundamental $SU(3)$ representation $1, m$. Thus in the case of $L \leq 3$, we have the following possibilities:

- $m = 3$. The only combination is $\{m_1, m_2, m_3\} = \{1, 1, 1\}$. Clearly this is an eigenstate of (1 2 3) with eigenvalue 1 and an eigenvector identified with the A_4 invariant singlet to be denoted $\underline{1}(3)$. The number in parenthesis indicates the m-value of the state.

- $\{m_1, m_2, m_3\} = \{0, 0, 0\}$. This leads to an A_4 invariant eigenstate denoted by $\underline{1}(0)$.

- $m = 2$. Now we have 1 possibility with three sets: $\{m_1, m_2, m_3\} = \{1, 1, 0\}, \{1, 0, 1\}, \{0, 1, 1\}$. Now the eigenvalues are ω, ω^2 and 1 with eigenvectors given as above. The transformation connecting the two basis given by:

$$\begin{pmatrix} \{1, 1, 0\} \\ \{1, 0, 1\} \\ \{0, 1, 1\} \end{pmatrix} = \frac{1}{\sqrt{3}}\begin{pmatrix} 1 & 1 & 1 \\ \omega^2 & \omega & 1 \\ \omega & \omega^2 & 1 \end{pmatrix}\begin{pmatrix} \underline{1}'(2) \\ \underline{1}''(2) \\ \underline{1}(2) \end{pmatrix} \tag{3}$$

- $m = 1$. Now we have two possibilities:

$$i)\{m_1, m_2, m_3\} = \{1, 0, 0\}, \{0, 1, 0\}, \{0, 0, 1\} \text{ and } ii)\{m_1, m_2, m_3\} = \{1, 1, -1\}, \{1, -1, 1\}, \{-1, 1, 1\}.$$

These two sets do not mix and again the eigenvalues are ω, ω^2 and 1. The transformation connecting the two basis given by:

$$\begin{pmatrix} \{1,0,0\} \\ \{0,1,0\} \\ \{0,0,1\} \end{pmatrix} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ \omega^2 & \omega & 1 \\ \omega & \omega^2 & 1 \end{pmatrix} \begin{pmatrix} \underline{1}_1^{'}(1) \\ \underline{1}_1^{''}(1) \\ \underline{1}_1(1) \end{pmatrix}, \tag{4}$$

$$\begin{pmatrix} \{1,1,-1\} \\ \{1,-1,1\} \\ \{-1,1,1\} \end{pmatrix} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ \omega^2 & \omega & 1 \\ \omega & \omega^2 & 1 \end{pmatrix} \begin{pmatrix} \underline{1}_2^{'}(1) \\ \underline{1}_2^{''}(1) \\ \underline{1}_2(1) \end{pmatrix}. \tag{5}$$

- $m = 0$. Now we have two possibilities:

  $i)\{m_1, m_2, m_3\} = \{1,-1,0\}, \{-1,0,1\}, \{0,1,-1\}$ and $ii)\{m_1, m_2, m_3\} = \{-1,1,0\}, \{1,0,-1\}, \{0,-1,1\}$.

  These two sets can be diagonalized separately. The transformation connecting the two basis given by:

$$\begin{pmatrix} \{1,-1,0\} \\ \{-1,0,1\} \\ \{0,1,-1\} \end{pmatrix} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ \omega^2 & \omega & 1 \\ \omega & \omega^2 & 1 \end{pmatrix} \begin{pmatrix} \underline{1}_1^{'}(0) \\ \underline{1}_1^{''}(0) \\ \underline{1}_1(0) \end{pmatrix}, \tag{6}$$

$$\begin{pmatrix} \{-1,1,0\} \\ \{1,0,-1\} \\ \{0,-1,1\} \end{pmatrix} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ \omega^2 & \omega & 1 \\ \omega & \omega^2 & 1 \end{pmatrix} \begin{pmatrix} \underline{1}^{'})_2(0) \\ \underline{1}_2^{''}(0) \\ \underline{1}_2(0) \end{pmatrix}. \tag{7}$$

There is no need to consider the negative M values, since their A_4 structure coincides with the above for $M \geq 0$. In the chosen basis the conjugate states coincide with these states for M even and they are opposite for M odd, a behavior typical of the angular momentum states. In any case the A_4 singlet states are specified by the corresponding M value for M=2, but in addition by a second a label a and b in the case of $M = \pm 1$ and $M = 0$.

If each particle is in the fundamental $SU(3)$ representation $1, m$ the possible three particle states are $[2\times1]^3$, $[2\times1]^2$, $[2\times1]^1$, $[0\times1]^1$, $[1\times1]^2$, $[1\times1]^1$ and $[1\times1]^0$, where the indicated numbers designate the angular momentum of the two particle state, the angular momentum of the third particle and, as an upper index, the total angular momentum of the thee particles respectively . The two particle states are designated in a similar manner $[1\times1]^L$ $L = 0, 1, 2$.

The $SU(3)$ states are $(\lambda, \mu) = (3,0), (1,1)$ and $(0,0)$, which correspond to the completely symmetric or $\underline{10}$-dimensional, the regular, $\underline{8}$ or octet, representation and the singlet completely antisymmetric representation. Their angular decomposition is as follows:

$$|(3,0), L = 3 \succ = [2\times1]^3, |(3,0), L = 1 \succ = -\frac{2}{3}[2\times1]^1 + \frac{\sqrt{5}}{3}[1\times1]^1, |(1,1)L = 1 \succ = \frac{2}{3}([0\times1]^1 + \frac{\sqrt{5}}{3}[2\times1]^1.$$

$$|(1,1)L = 2 \succ = ([2\times1]^2$$

It is now straightforward to expand the L, M states in terms of the three particle states$(1, m_1), (1, m_2), (1, m_3)$ via the Clebcsh-Gordan coefficients. Then transform them in the above A_4 basis to get the needed expansion of the $SU(3)$ states in terms of the singlet states of A_4. The results are:

1. For the $(\lambda, \mu) = (3,0) L = 3$ we get:

$$\begin{aligned} &(L = 3, M = \pm 3) \Leftrightarrow \underline{1}(3), \\ &(L = 3, M = \pm 2) \Leftrightarrow \underline{1}(2), \\ &(L = 3, M = \pm 1) \Leftrightarrow \frac{1}{\sqrt{5}} \left(2\underline{1}_1(1) + \underline{1}_2(1)\right), \\ &(L = 3, M = 0) \Leftrightarrow \sqrt{\frac{3}{10}} \left(\underline{1}_1(0) + \underline{1}_2(0)\right) + \frac{2}{\sqrt{10}}\underline{1}(0). \end{aligned} \tag{8}$$

   So this state is made completely out of invariant singlets $\underline{1}$

2. For the $(\lambda,\mu)=(3,0)L=1$ we find:

$$(L=1,M=\pm1)\Leftrightarrow-\frac{1}{9\sqrt{5}}\left(\underline{1}_1(1)-2\underline{1}_2(1)+10\left(\omega\underline{1}_1^{'}(1)-(1+\omega)\underline{1}_1^{''}(1)+\underline{1}'_2(1)+\underline{1}''_2(1)\right)\right),$$
$$(L=1,M=0)\Leftrightarrow\frac{1}{9\sqrt{5}}\left(10\left(\underline{1}_1^{'}(0)+\underline{1}_1^{''}(0)+\underline{1}'_2(0)+\underline{1}''_2(0)\right)+\underline{1}_1(0)+\underline{1}_2(0)\right)-\frac{1}{3\sqrt{15}}\underline{1}(0) \quad (9)$$

3. For the $(\lambda,\mu)=(1,1),L=1$ we get.

$$(L=1,M=\pm1)\Leftrightarrow\frac{1}{18}\left(\omega\underline{1}_1^{'}(1)+(1-\omega)\underline{1}_1''(1)+\underline{1}_1'(2)+\underline{1}_1''(2)+8\left(-\underline{1}_1+2\underline{1}_2\right)\right),$$
$$(L=1,M=0)\Leftrightarrow\frac{1}{18}\left(-\underline{1}_1^{'}(0)-\underline{1}_1^{''}(0)-\underline{1}_2^{'}(0)-\underline{1}_2^{''}(0)+8\left(\underline{1}_1(0)+\underline{1}_2(0)\right)\right)+\frac{4}{3\sqrt{3}}\underline{1}(0) \quad (10)$$

4. For the $(\lambda,\mu)=(1,1),L=2$ we have:

$$(L=2,M=\pm2)\Leftrightarrow\frac{1}{\sqrt{2}}\left(\underline{1}^{'}(1)+\underline{1}^{''}(1)\right),$$
$$(L=2,M=\pm1)\Leftrightarrow\frac{1}{2}\left(-\omega\underline{1}_1^{'}(1)+\omega\underline{1}_1^{''}(1)+\underline{1}_2^{'}(1)+\underline{1}_2^{''}(1)\right),$$
$$(L=2,M=0)=\frac{1}{2}\left(\underline{1}_1^{'}(0)-\underline{1}_1^{''}(0)-\underline{1}_2^{'}(0)+\underline{1}_2^{''}(0)\right) \quad (11)$$

5. For the the completely antisymmetric state $[1\otimes1]^1$ we have:

$$(L=0,M=0)=\frac{1}{\sqrt{2}}\left(\underline{1}_1(0)-\underline{1}_2(0)\right) \quad (12)$$

We summarize the results as follows.

- The A_4 singlets are characterized by an m quantum number. For $M\pm1$, 0 an additional label in needed, we used subscripts 1 and 2.
- The $L=3$ multiplet contains the A_4 invariant singlets only, so it stands is A_4 invariant. This is probably the reason for the claim that only this multiplet contains the A_4 singlet.
- The other $L=1,2$ multiplets may contain the invariant A_4 singlet, but only as a component together with the other A_4 singlets. If the symmetry is broken spontaneously so that the $\underline{1}_i^{''}(m)$ and $\underline{1}_i^{''}(m)$, $i=1,2$, acquire a vacuum expectation value as, e.g., in the context of a scalar potential, they will become A_4 invariant. This can happen since the quantum number L has nothing to do with angular momentum. It characterizes the $SU(3)\supset SO(3)$ flavor symmetry (see below), which, of course, is not sacred, but it must be somehow broken at some level.
- Representations with a number of quanta N, which is not a multiple of 3, cannot be represented solely in terms of A_4 singlets.Thus , e.g, for the $SU(3)$ representation (4,0) we have

$$(L=4,M=4)=\underline{1}(3),m_4=1;\ (L=4,M=3)=\frac{1}{2}\left(\underline{1}(3),m_4=0\right)+\frac{\sqrt{3}}{2}\left(\underline{1}(2),m_4=1\right)$$

$$((L=4,M=2)=\frac{1}{2\sqrt{7}}\left(\underline{1}(3),m_4=-1\right)+\frac{\sqrt{3}}{\sqrt{7}}\left(\underline{1}(2),m_4=0\right)+\frac{\sqrt{15}}{2\sqrt{7}}\left(\underline{1}(1),m_4=1\right)$$

$$((L=4,M=1)=\frac{\sqrt{3}}{2\sqrt{7}}\left(\underline{1}(2),m_4=-1\right)+\frac{\sqrt{15}}{2\sqrt{7}}\left(\underline{1}(1),m_4=0\right)+\frac{\sqrt{5}}{\sqrt{14}}\left(\underline{1}(0),m_4=1\right)$$

$$((L=4,M=0)=\frac{\sqrt{3}}{\sqrt{14}}\left(\underline{1}(1),m_4=-1\right)+\frac{\sqrt{4}}{\sqrt{7}}\left(\underline{1}(0),m_4=0\right)+\frac{\sqrt{3}}{\sqrt{14}}\left(\underline{1}(-1),m_4=1\right)$$

i.e. the magnetic substate m_4 enters lonely.

It is clear that any admissible polynomial of any member of the $L = 3$ multiplet is an A_4 invariant. Higgs potentials made of $SU(3)$ and $SO(3)$ invariants involving the other multiplets will have the proper structure (m is an additive quantum number and the singlets have the proper structure under conjugation). As we mentioned, when these non non invariant A_4 singlets acquire a vacuum expectation value, the symmetry will be broken down to A_4 via the scalar potential, which we will now discuss.

THE $SU(3)$ INVARIANT POTENTIAL

Constructing the $SO(3)$ invariants is well known.
The quadratic $SU(3)$ invariants are going to be of the form

$$[((\mu_1, \lambda_1), L) \otimes ((\lambda_1, \mu_1), L)]^{(00),0} \tag{13}$$

The quartic invariants are going to be of the form:

$$\Big[[((\lambda_1, \mu_1), L_1) \otimes ((\lambda_2, \mu_2), L_2)]^{((\lambda,\mu),L)} [((\lambda_3, \mu_3), L_3) \otimes ((\lambda_4, \mu_4)L_4)]^{((\mu,\lambda),L)}\Big]^{(0,0),0} \tag{14}$$

where the quantum number κ has been omitted for simplicity. The reduction $(0, \lambda) \times (\lambda, 0)$ is quite simple:

$$((0, \lambda), L_1)\,, \times ((\lambda, 0)L_2) = \sum_i ((\alpha_i, \alpha_i), L(i))$$

over all possible allowed combinations of the quantum numbers on the right. In the space of the above $SU(3)$ representations to construct the scalar potential up to quartic terms in these representations one needs the $SU(3)$ Wigner coefficients for the reductions:

i)

$$(3, 0) \times (0, 3) \to (3, 3) + (2, 2) + (1, 1) + (0, 0)$$

The last one is an SU(3) scalar, and can appear in the quadratic potential. All the resulting representations are self adjoined. So they can be coupled with theselves to yield a quartic $SU(3)$ scalar. The technology of the available $SU(3)$ coefficients [1] can easily be extended to evaluate all of them, as shown in the appendix.

ii) The self conjugate (octet) $SU(3)$ representation:

$$(1, 1) \times (1, 1) \to (3, 0) + (2, 2) + 2(1, 1) + (0, 3) + (0, 0)$$

The last one can appear in the quadratic term of the potential. The the available tables of $SU(3)$ coefficients [1] are adequate for their evaluation , but for the readers convenience we will supply them in the appendix. They can be appropriately combined to yield a quartic scalar.

iii) The coupling of the two considered representations, namely the decaplet and the octet

$$(3, 0) \times (1, 1) \to (4, 1) + (2, 2) + (1, 1) + (3, 0).$$

The result can be combined with the adjoined

$$(0, 3) \times (1, 1) \to (1, 4) + (2, 2) + (1, 1) + (0, 3)$$

to yield a quartic scalar. These, however, are not independent of those involved in i) and ii) above. In fact the resulting quartic coupling can be obtained by that combining i) and ii) via $SU(3)$ re-coupling coefficients. So we will not consider them further.

In the case i) the angular momentum decomposition is a follows

$$|(3,3)L\succ=\left\{\begin{array}{l}\kappa=1,L=\ 1\ \ 2\ \ 3\ \ 4\ \ 5\ \ 6\\ \kappa=3,L=\qquad 3\ \ 4\end{array}\right.,|(2,2)L\succ=\left\{\begin{array}{l}\kappa=0,L=\ 0\quad 2\quad\ 4\\ \kappa=2,L=\qquad 2\ \ 3\end{array}\right.,\ |(1,1)L=1,2\succ,\ |(0,0)L=0\succ$$

So one can construct 1 quadratic and 16 quartic $SU(3)\times SO(3)$ invariants. There exist 10 gauge bosons, to be eaten by the scalar non A_4 invariant components:

$$\underline{1}'_a(m),\underline{1}''_a(m),\ m=0,\pm1$$

In the case ii) above we have 3 possible combinations since there exist two possible $(\lambda,\mu)L=(1,1),L$ states. Then the angular momentum decomposition is:

$$|(2,2)L\succ=\left\{\begin{array}{l}\kappa=0,L=\ 0\quad 2\quad\ 4\\ \kappa=2,L=\qquad 2\ \ 3\end{array}\right.,\ |(1,1)L=1,2\succ_\rho,\ \rho=1,2,\ |(3,0)L=1,3\succ,\ |(0,3)L=1,3\succ,\ |(0,0)L=0\succ$$

(the index ρ appears since the octet can be obtained more than once). So one can construct 14 quartic $SU(3)\times SO(3)$ invariants. There exist 8 gauge bosons, to be eaten by the A_4 scalar components , e.g.

$$\underline{1}'_a(m),\ m=0,\pm1,\ \underline{1}'(m)(m),m=\pm2,\ \underline{1}''_a(m),\ m=0\pm1,\ \underline{1}^{''}_a(m),m=\pm2,a=1,2.$$

One can construct 10 more invariants by combining i) and ii) involving the self adjoined representations with the exception of (3,3) of i), without increasing the number of scalars.

Using the results obtained in the appendix one can write the invariant terms in terms of the component of the Higgs fields. Let us consider each case separately.

- The quadratic term (see Eq. 13). Then

$$[(\prec(\mu_1,\lambda_1),L)\otimes((\lambda_1,\mu_1),L)]^{(00),0}=\sum_M\prec(\mu_1,\lambda_1),L,-M,\ \lambda_1,\mu_1,LM|(0,0)0\succ$$
$$T^{(\mu_1,\lambda_1)},\kappa,L,-M\,T^{(\lambda_1,\mu_1},\kappa,L,M. \tag{15}$$

  where T indicate irreducible $SU(3)$ tensors. Using the conjugation properties of the SU(3) states [5], [6] one finds:

$$[(\prec(\mu_1,\lambda_1),L)\otimes((\lambda_1,\mu_1),L)]^{(00),0}=\frac{1}{\dim((\lambda_1,\mu_1))}\sum_{M\geq0}(2-\delta_{M0})\prec\lambda_1,\mu_1,L,M,|\lambda_1,\mu_1,L,M\succ. \tag{16}$$

  where the needed overlaps are computed via the expressions given above (see Eqs (8) and (9) for the decaplet and (10) and (11) for the octet).

- The quartic terns. Proceeding as above we find:

$$\Big[[((\lambda_1,\mu_1),L_1)\otimes((\lambda_2,\mu_2),L_2)]^{((\lambda,\mu),L)}\,[((\lambda_3,\mu_3),L_3)\otimes((\lambda_4,\mu_4)L_4)]^{((\mu,\lambda),L)}\Big]^{(0,0),0}=\frac{1}{\dim((\lambda,\mu))}$$
$$\sum_M\prec[((\lambda_1,\mu_1),L_1)\otimes((\lambda_2,\mu_2),L_2)]^{((\lambda,\mu),LM|)}\,|\,[((\lambda_3,\mu_3),L_3)\otimes((\lambda_4,\mu_4)L_4)]^{((\lambda),\mu,LM)} \tag{17}$$

  where the κ labels are understood. The needed overlaps are now complicated, but they can be computed using the C-G coefficients discussed in the appendix.

DISCUSSION

We have seen that by introducing only the $\underline{10}=(3,0)$ and $\underline{8}=(1,1)$ scalars of the group $SU_f(3)$ it is possible to construct more $SU_f(3)\supset SO(3)$ quadratic and quartic invariants than the number of non invariant A_4 singlets. So, in principle, possible to eat up all the $SU_f(3)$ gauge bosons after spontaneously breaking the symmetry. The needed $SU_f(3)\supset SO(3)$ C-G to this goal have been calculated. We have included the $\underline{10}=(3,0)$, because if its nice features regarding the invariantA_4 singlets it contains. Even if this representation is not contained in a higher symmetry group G, $G\supset SU_f(3)$, but the octet is, it may be adequate for the spontaneous symmetry breaking, since it contains invariant singlets . Furthermore the 14 invariants that one can construct from this alone is larger than the number of non invariant singlets contained in it. The needed $SU_f(3)\supset SO(3)$ C-G are provided in this work

APPENDIX: THE NEEDED $SU(3) \supset SO(3)$ C-G COEFFICIENTS

The full C-G coefficient needed for the reduction of $SU(3) \otimes SU(3)$ representations factorizes into one $SU(3) \supset SO(3)$ and one $SO(2) \supset SO(2)$ factor. Symbolically:

$$\prec (\lambda_1,\mu_1),\kappa_1,L_1,M_1,\ (\lambda_2,\mu_2,\kappa_2,L_2M_2|\lambda_3,\mu_3,\kappa_3,L_3M_3 \succ = \prec (\lambda_1,\mu_1),\kappa_1,L_1,\ (\lambda_2,\mu_2,\kappa_2,L_2||\lambda_3,\mu_3,\kappa_3,L_3 \succ$$

$$\prec L_1M_1,L_2,M_2|L_3,M_3 \succ .$$

So, since the usual $SO(2) \supset SO(2)$ C-G coefficient is well known, we need consider only the double bar coefficients. The only $SU(3) \supset SO(3)$ C-G coefficients, not found in the literature [1] are those involving the $(3,0) \otimes (0,3)$ reduction. On this occasion we will consider the general reduction

$$(\lambda,0) \otimes (0,3) \to (\lambda',\mu'),\ (\lambda',\mu') = (\lambda,3),\ (\lambda-1,2),\ (\lambda-2,1),\ (\lambda-3,0)$$

To this end we will use the build-up procedure [1]

$$\prec (\lambda,0)L,(0,3)\ell||(\lambda',\mu')\kappa'L' \succ U\left((\lambda,0),(0,2),(\lambda',\mu')(0,1)1;(\lambda^{''},\mu''),(0,3)\right) =$$

$$\sum_{\kappa''L''\ell_1} U(L,\ell_1,L',1;L^{''}\ell) \prec (\lambda,0)L,(0,2)\ell_1||(\lambda^{''},\mu'')\kappa^{''}L^{''} \succ$$

$$\prec (0,2)\ell_1,(01)1||(03)\ell \succ \prec (\lambda^{''},\mu'')\kappa^{''}L^{''},(0,1)||(\lambda',\mu')\kappa'L' \succ$$

where $U(L,\ell_1,L',1;L^{''}\ell)$ is the (unitary) Racah 6j function for the rotation group and $U\left((\lambda,0),(0,2),(\lambda',\mu')(0,1);(\lambda^{''},\mu''),(0,3)\right)$ is an analogous function for the SU(3) group, which can be judiciously chosen to make the above expression as simple as possible. It can be computed by requiring the C-G coefficients of the left hand side to be suitably normalized. A convenient choice is

$$(\lambda^{''},\mu'') = (\lambda,2),\ (\lambda-1,1),\ (\lambda-2,0),\ (\lambda-2,0) \Leftrightarrow (\lambda^{'},\mu') = (\lambda,3),\ (\lambda-1,2),\ (\lambda-2,1),\ (\lambda-3,0).$$

Using Eq. (25a) of the above reference it can be cast in the form:

$$\begin{aligned}\prec (\lambda,0)L,(0,3)\ell||(\lambda',\mu')\kappa'L' \succ \quad & U\left((\lambda,0),(0,2),(\lambda',\mu')(0,1);(\lambda^{''},\mu''),(0,3)\right) = (-1)^{L-L'+(2-\ell_1)/2}(-1)^{(\mu'-\lambda'+\lambda-3)/3} \\ & \sqrt{\frac{\dim(\lambda',\mu')}{\dim(\lambda,0)}}\sqrt{\frac{2L+1}{2L'+1}}\sum_{\kappa''L''\ell_1}\sqrt{\frac{5}{3}\frac{2\ell_1+1}{2\ell+1}}U(L,\ell_1,L',1;L^{''}\ell) \\ & \prec (\lambda^{''},\mu'')\kappa^{''}L^{''},(2,0)\ell_1||(\lambda,0)L \succ \\ & \prec (0,3)\ell,(1,0)1||(02)\ell_1 \succ \prec (\lambda',\mu')\kappa'L',(1,0)1||(\lambda^{''},\mu'')\kappa^{''}L^{''} \succ \end{aligned} \tag{18}$$

with $\dim(\lambda,\mu) = \frac{1}{2}(\lambda+1)(\mu+1)(\lambda+\mu+2)$ the dimension of (λ,μ). All the $SU(3)$ G-C entering the right hand side of the previous equation can be found in the tables [1]. The parameter κ needed in case there exists degeneracy in L, will be omitted in the cases that such a degeneracy is not present. We have obtained analytic expressions for the results, but their quite cumbersome. So we will present here the results needed here. We should mention that only the entries with non zero valus of the C-G are included in the tables below.

$$(3,0)L,(03)\ell||(0,0)L' = 0 \succ = \begin{cases} \sqrt{\frac{3}{10}} & L=\ell=1 \\ -\sqrt{\frac{7}{10}} & L=\ell=3 \end{cases} ; (3,0)L,(03)\ell||(1,1)L' = 1 \succ = \begin{cases} L & \ell & C-G \\ 1 & 1 & \sqrt{\frac{14}{15}} \\ 3 & 3 & -\sqrt{\frac{1}{15}} \end{cases}$$

$$(3,0)L,(03)\ell||(1,1)L'=2\succ=\begin{Bmatrix} L & \ell & C-G \\ 1 & 1 & -\frac{3}{5}\sqrt{\frac{3}{5}} \\ 1 & 3 & \frac{2}{5}\sqrt{\frac{7}{5}} \\ 3 & 1 & \frac{2}{5}\sqrt{\frac{7}{5}} \\ 3 & 3 & -\frac{1}{5}\sqrt{\frac{6}{5}} \end{Bmatrix};3,0)L,(03)\ell||(2,2)\kappa'=0,L'=4\succ=\begin{Bmatrix} L & \ell & C-G \\ 3 & 3 & -\sqrt{\frac{11}{35}} \\ 1 & 3 & 2\sqrt{\frac{3}{35}} \\ 3 & 1 & -2\sqrt{\frac{3}{35}} \end{Bmatrix}$$

$$(3,0)L,(03)\ell||(2,2)\kappa'=0,L'=2\succ=\begin{Bmatrix} L & \ell & C-G \\ 3 & 3 & \frac{12}{5}\sqrt{\frac{2}{91}} \\ 1 & 3 & -\frac{1}{5}\sqrt{\frac{39}{7}} \\ 3 & 1 & \frac{22}{5}\sqrt{\frac{3}{91}} \\ 1 & 1 & \frac{2}{5}\sqrt{\frac{1}{3}} \end{Bmatrix};(3,0)L,(03)\ell||(2,2)\kappa'=2,L'=2\succ=\begin{Bmatrix} L & \ell & C-G \\ 3 & 3 & -\frac{12}{5}\sqrt{\frac{6}{65}} \\ 1 & 3 & -\frac{2}{5}\sqrt{\frac{13}{5}} \\ 3 & 1 & -\frac{2}{5}\sqrt{\frac{21}{65}} \\ 1 & 1 & \frac{1}{5}\sqrt{\frac{1}{65}} \end{Bmatrix}$$

$$(3,0)L,(03)\ell||(2,2)\kappa'=2,L'=3\succ=\begin{Bmatrix} L & \ell & C-G \\ 3 & 3 & 3\sqrt{\frac{3}{35}} \\ 1 & 3 & -2\sqrt{\frac{1}{35}} \\ 3 & 1 & 2\sqrt{\frac{1}{35}} \end{Bmatrix};(3,0)L,(03)\ell||(2,2)\kappa'=0,L'=0\succ=\begin{Bmatrix} L & \ell & C-G \\ 3 & 3 & -\sqrt{\frac{3}{10}} \\ 1 & 1 & -\sqrt{\frac{7}{10}} \end{Bmatrix}$$

$$(3,0)L,(03)\ell||(3,3)\kappa'=1,L'=1\succ=\begin{Bmatrix} L & \ell & C-G \\ 1 & 1 & -\sqrt{\frac{14}{15}} \\ 3 & 3 & -\sqrt{\frac{1}{15}} \end{Bmatrix};(3,0)L,(03)\ell||(3,3)\kappa'=1,L'=2\succ=\begin{Bmatrix} L & \ell & C-G \\ 1 & 1 & \frac{2\sqrt{2}}{5} \\ 1 & 3 & -\frac{2}{5}\sqrt{\frac{6}{5}} \\ 3 & 1 & \frac{2}{5}\sqrt{\frac{6}{5}} \\ 3 & 3 & \frac{1}{5}\sqrt{\frac{1}{7}} \end{Bmatrix}$$

$$(3,0)L,(03)\ell||(3,3)\kappa'=1,L'=3\succ=\begin{Bmatrix} L & \ell & C-G \\ 1 & 3 & -\sqrt{\frac{3}{55}} \\ 3 & 1 & -4\sqrt{\frac{3}{55}} \\ 3 & 3 & 2\sqrt{\frac{1}{55}} \end{Bmatrix};(3,0)L,(03)\ell||(3,3)\kappa'=3,L'=3\succ=\begin{Bmatrix} L & \ell & C-G \\ 1 & 3 & -8\sqrt{\frac{1}{77}} \\ 3 & 1 & 1\sqrt{\frac{1}{77}} \\ 3 & 3 & -2\sqrt{\frac{3}{77}} \end{Bmatrix}$$

$$(3,0)L,(03)\ell||(3,3)\kappa'=1,L'=4\succ=\begin{Bmatrix} L & \ell & C-G \\ 1 & 3 & 3\sqrt{\frac{11}{287}} \\ 3 & 1 & 4\sqrt{\frac{11}{287}} \\ 3 & 3 & -2\sqrt{\frac{3}{287}} \end{Bmatrix};(3,0)L,(03)\ell||(3,3)\kappa'=3,L'=\succ=\begin{Bmatrix} L & \ell & C-G \\ 1 & 3 & 8\sqrt{\frac{1}{205}} \\ 3 & 1 & -3\sqrt{\frac{1}{205}} \\ 3 & 3 & 2\sqrt{\frac{33}{205}} \end{Bmatrix}$$

$$(3,0)L,(03)\ell||(3,3)\kappa'=1,L'=5\succ=-1\,;(3,0)L,(03)\ell||(3,3)\kappa'=1,L'=6\succ=1$$

Incidentally the $SU(3)$ U-functions are:

$$U((30)(02)(\lambda',\mu')(01);(\lambda'',\mu'')(03)=1,\frac{1}{\sqrt{2}},\frac{\sqrt{7}}{3},1$$

for

$$(\lambda',\mu'),(\lambda^{''},\mu^{''})=(00),(1,0);\ (1,1),(1,0);\ (2,2)(2,1);\ (3,3)(3,2)$$

respectively.

The needed $SU(3)$ C-G involving the representation $\underline{8}=(11)$ can be obtained from the existing tables [1], but, for the reader's convenience, we will include them here.

$$(1,1)L,(11)\ell||(0,0)L'=0\succ=\begin{cases} -\sqrt{\frac{3}{8}} & L=\ell=1\\ \sqrt{\frac{5}{8}} & L=\ell=2\end{cases}$$

$$(1,1)L,(11)\ell||(1,1),L'=1\succ_1=\begin{cases} L & \ell & C-G\\ 1 & 1 & -\sqrt{\frac{1}{6}}\\ 3 & 3 & \sqrt{\frac{5}{6}}\end{cases};(1,1)L,(1,1)\ell||(1,1)L'=1\succ_2=\begin{cases} L & \ell & C-G\\ 2 & 1 & \sqrt{\frac{1}{2}}\\ 1 & 2 & \sqrt{\frac{1}{2}}\end{cases}$$

$$(1,1)L,(11)\ell||(1,1),L'=2\succ_1=\begin{cases} L & \ell & C-G\\ 2 & 1 & \sqrt{\frac{1}{2}}\\ 1 & 2 & -\sqrt{\frac{1}{2}}\end{cases};(1,1)L,(1,1)\ell||(1,1)L'=2\succ_2=\begin{cases} L & \ell & C-G\\ 1 & 1 & -\sqrt{\frac{3}{10}}\\ 2 & 2 & \sqrt{\frac{7}{10}}\end{cases}$$

$$(1,1)L,(11)\ell||(3,0),L'=3\succ_1=\begin{cases} L & \ell & C-G\\ 2 & 1 & -\sqrt{\frac{1}{4}}\\ 1 & 2 & \sqrt{\frac{1}{4}}\\ 2 & 2 & \sqrt{\frac{1}{2}}\end{cases};(1,1)L,(1,1)\ell||(3,0)L'=1\succ_2=\begin{cases} L & \ell & C-G\\ 2 & 1 & \sqrt{\frac{1}{2}}\\ 1 & 2 & \sqrt{\frac{1}{2}}\end{cases}$$

$$(1,1)L,(11)\ell||(0,3),L'=3\succ_1=\begin{cases} L & \ell & C-G\\ 2 & 1 & -\sqrt{\frac{1}{4}}\\ 1 & 2 & \sqrt{\frac{1}{4}}\\ 2 & 2 & -\sqrt{\frac{1}{2}}\end{cases};(1,1)L,(1,1)\ell||(0,3)L'=1\succ_2=\begin{cases} L & \ell & C-G\\ 2 & 1 & \sqrt{\frac{1}{2}}\\ 1 & 2 & -\sqrt{\frac{1}{2}}\end{cases}$$

$$(1,1)L,(11)\ell||(2,2),L'=0\succ=\begin{cases} L & \ell & C-G\\ 1 & 1 & -\sqrt{\frac{5}{8}}\\ 2 & 2 & \sqrt{\frac{3}{8}}\end{cases};(1,1)L,(1,1)\ell||(2,2)L'=3\succ=\begin{cases} L & \ell & C-G\\ 2 & 1 & \sqrt{\frac{1}{2}}\\ 1 & 2 & \sqrt{\frac{1}{2}}\end{cases}$$

$$(1,1)L,(11)\ell||(2,2),\kappa'=0,L'=2\succ=\begin{cases} L & \ell & C-G\\ 1 & 1 & -\frac{1}{2}\sqrt{\frac{7}{13}}\\ 2 & 1 & -\frac{1}{2}\sqrt{\frac{21}{13}}\\ 1 & 2 & \frac{1}{2}\sqrt{\frac{21}{13}}\\ 2 & 2 & \frac{1}{2}\sqrt{\frac{3}{13}}\end{cases};(1,1)L,(1,1)\ell||(2,2)\kappa'=2,L'=2\succ=\begin{cases} L & \ell & C-G\\ 1 & 1 & \frac{7}{2}\sqrt{\frac{3}{65}}\\ 2 & 1 & -\frac{1}{2}\sqrt{\frac{5}{13}}\\ 1 & 2 & \frac{1}{2}\sqrt{\frac{5}{13}}\\ 2 & 2 & -\frac{3}{2}\sqrt{\frac{7}{65}}\end{cases}$$

$$(1,1)L,(11)\ell||(2,2),L'=4\succ=4$$

Acknowledgements: The author is indebted to Professors G. K. Leontaris for useful discussions and Tony Thomas for his hospitality and support during his visit in Adelaide.

[1] J.D. Vergados, Nuc. Phys. A, **111** (1968) 681.
[2] J. Berger and Y. Grossman, JHEP, **1002**(2010) 071.
[3] B. A. Ovrut, J. Mah. Phys. **19**(1978) 418,doi:10.1063/1.523660.
[4] G. Elesi, J. Mah. Phys. **37 (4)**(1996) 1596,doi:10.1063/1.531470.
[5] J. J. de Swart, Rev. Mod. Phys. **35** (1963) 916
[6] J. P. Elliott and M. Harvey, Proc. Roy. Soc. **A 272** (1963) 557.